

Humans In Variety / Adapted Immune Developmental Symbiosis

By Christl J. Meyer, Biologist, Vienna/ Austria, 2010-06-27

Abstract/ Summary

In this paper I aim demonstrate a highly novel insight into the evolution of the Immune System of Humans. The Human Genome Project which mapped out our genetic background by reducing the amount of protein coding genes to approximately 20,000 and augmented the RNA transcriptions, assumes vast proportions for future research which is centred on health and disease.

Thus the MHC-complex and the HLA-system provide new insights into the co-evolution of the species particularly of mammals (and primates) with microbes. We have to take in account that evolution is an ongoing process which renders creatures susceptible to diseases in connection with environmental stimuli –specifically stress- leading to insights in the causes of many complex diseases like malaria, diabetes, lupus, multiple sclerosis, rheumatoid arthritis, many infections, influenza and cancer.

The HIV/AIDS- hypothesis will be dismantled, specifically, the notion of the sexual transmission of a virus. Instead I propose that an evolutionary adaption of an ancient retrovirus established in the human genome renders viruses in a symbiotic interaction with the host genes of the immune system to participate in cell communication.

The ethics of the 21st century makes us responsible for not prolonging an unwanted genocide by underestimating weaker voices in science. Instead we have to focus on the latest research and apply the results to an overhaul of our understanding of AIDS.

- I. Introduction: New Scientific Projects
- II. Genetic Diversity of Humans
- III. The MHC/HLA-System and its Origin
- IV. Sperm Proteins, Pregnancy and Protection of Health
- V. Auto-, Allereactions and Diseases
- VI. Evolution Never Ends
- VII. About Testing, Responsibility and Ethics

I. Introduction

As previously shown in my paper “Reconciliation between Pure Scientists and AIDS-Dissidents:

Could an ancient retrovirus, RNA-interference and stress be the answer to the divergent opinions? “ there is growing evidence, that HIV is an ancient retrovirus acting as genes that can be influenced by environmental stress and small RNAs.

The Human Genome Project (HGP) provided information about less protein coding genes than we were previously aware. Instead we are now eagerly discovering the Human Epigenome (HEP), looking for the methylated genes [1,4] and according to Jenuwein and Allis [2] histones of the DNA because of the great importance of gene expression in the different cells and tissues of our bodies[3].

The Human Microbiome Project (HMiP) aims to specify the genes of the microbes living on and in our bodies [5]. A single human has 10fold more microbes than cells and the genes of our symbioses, which contribute to our digestion, vitamin supplementation and gene activation are now of greater interest to research. The co-evolution of bacteria and their viruses provides new information about the acceleration of evolution [6, 8] and lateral (horizontal) gene transfer from microbes to their hosts [7]. The Human Protein Project (HPP) and the Human Metabolome Project (HMP) will provide additional information on the complexity of life and evolution.

II. The Diversity of Humans

“The genetic structure of the indigenous hunter-gatherer peoples of southern Africa, the oldest known lineage of modern humans, is important for understanding human diversity “[9]. These hunter-gatherers, known as Khoisan, San, or Bushmen, are genetically divergent from other humans. In terms of nucleotide substitutions, the Bushmen seem, on average, to differ more from each other than, for example, a European or an Asian. There is also a discontinuity between local hunter-gatherers and central Europe’s first farmers [10]. By analyzing ancient DNA of Neolithic hunter-gatherers and contemporary Scandinavians, Malstrom et al. revealed a lack of continuity [11]. Humans underwent an adaptation process that was influenced by geography [12]. This resulted in differences including in the immune system. People of African descent show reduced neutrophil count due to a regulatory variant [13]. There is also an extensive genetic diversity in the HLA class II region of Africans from Gambia and Malawi. This diversity is twice as extensive as found in northern Europeans [14].

Of consequence, we find differences in humoral responses between Ethiopian and Swedish people who are claimed to be “infected” by HIV [15]. Therefore it is necessary to investigate the nature and biological background of the immune system responsible for cell activation, receptor and antibody generation and the communication between all partners involved in the immune response.

III. The MHC/HLA-System and its Origin

A most interesting study by Mungall et al. reports that chromosome 6 constitutes about 6% of the human genome and harbours 1,557 genes and 633 pseudo-genes [16, 30]. Within the essential immune loci of the major histocompatibility complex, HLA-B was found to be the most polymorphic gene. Among these are genes directly implicated in diseases such as cancer and autoimmunity. Having a look at the supplementary tables, which give full lists of HLA allele-associated HIV polymorphisms in Protease, Reverse Transcriptase, VPR and Nef, they show that more than 240 gene variants of Nef are adapted to HLA. There are also tremendous variants for the other genes. The main effect of Nef is to block transport of MHC-I molecules to the cell surface, leading to accumulation in intracellular organelles [23a].

This study and the report from Brumme et al. [25] confirms the nature of HIV – showing that it is of endogenous origin – part of our genome and constituted by the high variability of the genes, which elucidates the variability of the HIV-variants and the immune escape. As Hedrick stated previously in 1994, there is a high heterozygosity, reaching over 60% at some amino acid sites with primary function. In some populations there is an observed deficiency in homozygotes. There seems to be a balancing selection in the MHC region related to the function for protection against microbes. Probably there is a selection in the MHC involved in non-random mating and maternal-fetal interactions in pregnancy [17].

The Mexican cohort [27] and the ANRS Genome Wide Association 01 Study [28] conclude that HLA/MHC controls the HIV-reservoir and replication.

Gyllenstein et al. reported an allelic diversity that is generated by intraexon sequence exchange at the DRB1 locus of primates [18]. Doxiadis et al. report phylogenetic evidence that supports the notion of the generation of new HLA-DRB genes as a dynamic and steadily ongoing process. This is due to the presence of indels (insertions/deletions), mainly mapping to introns. The research compared a large number of full-length sequences of rhesus macaques, chimpanzees and humans. As no evidence was found for convergent evolution, the combination of these observations indicates that ancient peptide binding motifs are frequently reshuffled among duplicated members of the HLA-DRB multigene family [19]. Oosterhout stated in his review on population genomics and epidemiology the concern of transposons in the MHC of the vertebrate immune system. Transposons constitute a large proportion of the vertebrate genome, and on average more than 40% of the mammalian genome consists of these parasitic elements. TEs have a non-random distribution throughout the genome, and they show an increased density in the MHC in a wide range of vertebrates [20].

Genetic drift may play an important role in the population genetics as Moore et al. argue [21]. Characteristic mutations in HIV RT are evident at the population level. For-example, the presence of HLA-B7 in the Western Australian HIV Cohort Study is strongly associated with non-synonymous mutation at position 135 of HIV RT which is an anchor residue of a HLA B51 restricted CTL epitope. This mutation allows for escaping the host CTL response. These results are important because of the occurrence of drug- induced mutations [21].

Among the many host cell-derived proteins found in HIV-1, HLA-II appears to be selectively incorporated onto virions. [21a]. The genetic diversity of the envelope glycoprotein from HIV-1 isolates from eight countries in Africa is studied by Louwagie et al. The data confirm the existence of several genetic subtypes and broaden the genetic variability observed for envelope subtypes. The geographic spread of different subtypes was shown to be substantial [21b]. “Rapid Evolution of Major Histocompatibility Complex Class I Genes in Primates Generates New Disease Alleles in Humans via Hitchhiking Diversity” The diversity created by single nucleotide variations (SNV) was not evenly distributed. It was rather concentrated within the gene-clusters HLA-A and HLA-B/C. These polymorphisms seem to be species specific. They might have been selected in adaption to the constantly evolving microbial

antigenic repertoire [26]. The Mexican cohort [27] and the ANRS Genome Wide Association 01 Study [28] found that HLA/MHC controls HIV-reservoir and replication. Clerici and Shearer [22] present a model “The Th1-Th2 hypothesis of HIV infection: new insights” where they claim activation-induced, cytokine- modulated, programmed cell death as a major factor in the pathogenesis of HIV infection in AIDS. Immunoregulatory cytokines are also produced by non T-cells, including monocytes/macrophages, natural killer cells and B-cells. So the authors prefer the terms type 1 and type 2 responses. They suggest an endogenous imbalance in the immunoregulatory cytokine network. HIV resistance in female sex-workers in Northern Thailand seems to be influenced by synergistic impact of HIV-specific cytotoxic T lymphocytes, HLA-A11, and chemokine –related factors [23]. Transcriptional analysis for host factors required by HIV-1 was performed by RNA interference. More than 250 HIV-dependency factors have been identified. These proteins participate in cellular functions. Transcriptional analysis revealed that these genes were enriched for high expression in immune cells [24]. In addition HIV incorporates HLA-DR, which is a cell-surface protein in enormous quantities. Other proteins that have been found are HLA class I proteins and various cell adhesion proteins as well as proteins from inside the cell like cyclophilin A, actin and ubiquitin [29]. Thus HIV has not been proven to be a virus. **Conclusion: HIV is an evolutionary adapted and partly active variable and heritable gene construct in our immune (MHC/HLA)-system.**

IV. Sperm Proteins, Pregnancy and Protection of Health

The influence of human semen on immunity is of importance for estimating the impact on reactions concerning sexuality and child development. Jeremias et al. claim that human semen is both an inducer of an anti-inflammatory TH2 immune response and an inhibitor of TH1 cell mediated immunity. The induction of interleukin 10 and 70 kDa heat shock protein gene transcription and IFN- γ was examined [31]. Virus-encoded (ORF) homologs of cellular interleukin-10 range in sizes for cellular Il-10 proteins [32]. The result in both cases is a shift from a T1 to T2 immune response which protects cells from immune attack, being an absolute necessary prevention for the foetus, which could be otherwise attacked by maternal response to paternal antigens in the womb. There is a cross-reactivity of sperm and T-lymphocyte antigens that results in higher titers of antibodies in couples with antisperm immunity as compared with “normal” couples [33]. A number of studies have suggested that an immune response to human leukocyte antigen HLA alloantigens may contribute to protection against HIV infection [34]. Infectivity for HIV through heterosexual transmission is low, and sexual transmitted diseases (STDs) may be the most important cofactor for transmission [35]. Heterosexual and homosexual monogamous partners practising unprotected sex develop CD4+ and CD8+T cell proliferative responses to the partners’ unmatched cells and a minority may be tolerated. These together with other findings suggest that allogeneic immunity may play a significant role in HIV [36]. HIV-1 gp120 is an immunoglobulin superantigen, which can bind to pre-immune serum Ig. The level of pre-immune anti-gp120 IgG is a polymorphic population trait, and low levels are a potentially specific and significant factor in homosexual transmission of HIV infection [37]. HIV-1 binds and enters normal sperm and can transfer HIV-1 like particles to normal human oocytes [38]. In 1991 Calarco and Whitmer proposed that HIV-1 expression could occur during early mammalian development [39]. Maternal (rather than paternal) allergy background confers stronger allergy risk for the offspring. Maternal responses to foetal antigens were related to foetal immune responses and

subsequent allergy. The number of previous pregnancies was associated with stronger maternal responses to foetal alloantigens [40].

Conclusion: HIV is a natural product in sperm which has a relation to HLA and protects the foetus from maternal rejection of paternal antigens by shifting T1 to T2. Heterosexual transmission of HIV is only suggested with additional pathogens in STDs. Homosexual transmission is due to rejection of alloantigens. Allogeneic immunity protects from infection but can be related to allergies also in the offspring.

V. Auto-, Allereactions and Diseases

Concerning the mentioned HIV is a regulatory and even life- promoting element of the immune system which has evolved over millions of years as a symbiotic partner that interacts in health and disease. Multiple interactions in cell communication have been proven concerning HIV specifically in GALT (gut associated lymphoid tissue), which makes sense for protecting the body from foreign invaders. Mehandru et al. emphasize that the gastrointestinal tract -associated lymphoid tissue constitutes the largest immune compartment in the body. More than 60% of the bodies' total lymphocytes are estimated to be T-cell associated with the small intestinal epithelium [45]. Dissemination of a virus to GALT is mediated by an integrin and Gp 120 leading to the formation of virological synapses, which facilitate efficient cell-to-cell spreading of HIV-1 [41]. Retroviral assembly is driven by Gag release, which is promoted by the clathrin adaptor complex AP-1 to intracellular sites of active budding – the machinery that forms intraluminal vesicles of the multivesicular body MVB. Protein sorting is critical for diverse cellular functions, such as receptor down-regulation, degradation of membrane proteins and lysosome like organelles, which includes attachment of ubiquitin to cargo proteins [42]. Teis et al. state that a certain complex (ESCRT) is required for cargo sequestration and vesicle formation during MVB sorting [43]. In addition direct cell-cell communication mediated by plasma membrane-spanning gap junction (GJ) channels is vital to all aspects of cellular life. Cells internalize GJ in response to various stimuli. In this process, clathrin, dynamin (GTPase) and other proteins are involved in internalizing double-membrane vesicles into cells [44]. Exosomes correspond to the multivesicular body and are released upon exocytic fusion with the plasma membrane. They function in intercellular communication during the immune response. They might be involved in tissue developmental processes and seem to be of ancient origin [46]. Stephen J. Gould proposed the “Trojan exosome hypothesis” in which retroviruses exploit a cell-encoded pathway of intercellular vesicle traffic, exosome exchange and last but not least that alloimmunity is a central component of antiretroviral immunity [47]. Antibodies against HLA neutralize HIV-1 in vitro. This was demonstrated in alloimmune sera from polytransfused patients [50]. Frank P. Ryan refers to “mutualistic symbiosis and HERVs in our genomes that have lost the ability to survive independently, but their removal from our genome would also make us extinct” [48].

The American Society of Microbiology regards the subject of viral contribution to host evolution as so vital that it commissioned Luis Villareal to write a book on this issue to educate the next generation of scientists (Viruses and the evolution of life) [49]. The envelope glycoprotein of HIV-1 gp120 has been identified as a member of the Immunglobulin superantigens (Ig-SAg) that selectively binds to an unusually high proportion of **endogenous nonimmune Ig** that are members of the VH3 Ig gene family [51]. The importance for diseases might be that the up-regulation of expression of endogenous retroviral superantigens has substantial implications for understanding the pathology of virus infections i.e. Epstein-Barr virus [52]. SAgS seem to be involved in allergy and autoimmune diseases [53]. Thus they

might be used as a therapeutic agent in the treatment of cancer. The light chain subunits of antibodies cloned from patients with systemic lupus erythematosus bind and hydrolyze gp120 sAg [55]. Already in 1990 studies on other lupus patients reported that about one-third of them produced antibodies to the p24 gag protein of HIV-1 as demonstrated in Western blotting [56]. In a study of HLA class I and II antigens in South African Blacks with Grave's disease, there was a significant increase in the frequency of HLA-DR3 in patients compared to control subjects, and a relationship in the DRI locus [54]. Grave's disease is an autoimmune disorder. A study of multiple sclerosis patients provides "solid evidence" that HTLV-I, which is similar to HIV, is involved in the MS disease process [57]. Approximately 25% of severe haemophilia A (HA) patients develop antibodies to factor VIII protein, which is due to impact of polymorphisms of the MHC complex class II and other factors like interleukin-10 [58]. As antibodies to blood products were defined as HIV contaminated in previous studies [59] the reported data from 2009 may now be interpreted as gene expression of distinct SNPs.

There is also a relation of HIV-1 acquisition to hormonal contraception and to herpes simplex virus type 2 among Kenyan women [60]. The genetic predisposition to type 1 -diabetes is associated with genes of the HLA system, specifically with HLA-DR and -DQ [61]. Research on transgenic mice resulted in activation of gene expression in HIV by Mycobacterium tuberculosis and suppression after antimycobacterial chemotherapy [62]. This shows the bystander function of HIV in tuberculosis. As a result, HIV is not the cause of the disease but tuberculosis is the disease and HIV is part of the communication system and part of an active immune system. Gene regulation is subject to hormone control specifically to corticosteroids in retroviral systems and to pregnancy in women [63]. This might indicate stress and placental involvement of testing HIV positive. The HIV-1 VPR-protein might be protective against cancer by inducing apoptosis in tumour cells [64]. Cell surface MHC class I-like proteins are up-regulated upon cell stress, including viral and bacterial infection and tumour transformation and are recognized by NKG2D a C-type lectin- activating receptor [65]. Stress and depressive symptoms are associated with decrease of protective NK and CD8+ T lymphocytes in HIV-infected men [66]. Individuals from Central Africa have a higher level of immune system activation compared with non-African populations which might be due to multiple and frequent exposures to viral, bacterial and parasitic antigens [67].

By comparing medical treatment for "HIV-infection" resistance to zidovudine was significantly higher in individuals with disease progression than in those from the control group [68]. The variable region (V3) of the gp 120 surface envelope glycoprotein of HIV-1 is a highly variable disulfide-bonded structure which triggers cell infection and escape from antiviral drugs, specifically entry inhibitors and is a target for neutralizing antibodies [69]. After effective Highly Active Antiretroviral Therapy (HAART) people with HIV might experience an "immune restoration syndrome" which is established by lymphocyte recovery period and might be manifested by infectious agents such as cytomegalovirus or mycobacterium avium intracellulare or a sudden onset of sarcoidal granulomatous reactions. An uncontrolled Th1 response as a result of cytokine alterations via Il-2 is the causative mechanism [70] as the authors of this study claim. In "HIV infection" Il-1 is increased which shifts the reactions to a Th2 response. Reuse, Calao Kabeya et al. from Belgium propose a synergistic activation of HIV-1 expression by deacetylase inhibitors and prostratin as implications for treatment of latent infections. This might reduce the size of latent HIV-1 reservoirs in HAART-treated patients [71]. A study from 2008 found an inactivation of HIV-1 by modification of nucleocapsid zinc fingers [72].

Conclusion: HIV is a symbiotic agent in GALT and other tissues, thus protective against microbes due to nutrition. Exosomes are produced and function in cell communication processes. Gp 120 is active as a superantigen that increases Th2 related antibody production in infections as a "booster" and might be due to allergy and autoimmunity.

Stress is involved in gene expression. HIV might be protective in some cancers. Medications and HAART might have different (negative) impacts on the balance of the Th1 / Th2- system.

VI. Evolution Never Ends

Human endogenous retroviruses (HERVs) have been estimated to be part of the genome and are replication incompetent. HERV-W encodes a highly fusogenic membrane glycoprotein within functional retrovirus genes which has been proposed to play a role in normal placental development not only in humans but also in simian and pig cells. The HERV-W entered the genome of primates approximately 25 million years ago. They are replication defective because of mutations within functional retrovirus genes. The existence of individual open reading frames corresponding to gag, pol, and env have been shown to encode proteins in some cases. HERVs could be potentially assembled into infectious virions through transcomplementation with virion proteins encoded by different HERVs [73].

This research from 2000, by Dong Sung An et al. from the UCLA AIDS Institute was done before the Human Genome Project was published and the tremendous occurrence of HERVs were detected in the Human genome specifically in relation to the MHC/HLA. The authors also claim that a functional envelope glycoprotein would confer the ability to be transmitted vertically and / or horizontally. Kumar et al. revealed a two fold higher expression of CXCR4 mRNA in early as compared to term human placenta. Chemokines and their receptors may play a crucial role in angiogenesis and proliferation in cell function. The receptor expression may be developmentally regulated and its role in the early stages of pregnancy is implicated when embryogenesis and organogenesis occurs[74]. They suggest that CXCR4 may not have a direct role in HIV infection, as only 1-2% of the placental transmission of HIV takes place in the early placenta. Allogeneic stimulation in early pregnancy improves pre- and postnatal ontogenesis by activation of the female immune system and enhancement of rise of plasma progesterone. The dissimilarity of mother and foetus induces stress resistance in the progeny of BALB/cLac mice [75]. This evolutionary development acts also as a mechanism for creating phenotypic diversity [77]. In seropositive children, a well- known cross-reactivity between HLA-DR and gp 120 is marked [76], [78].

Conclusion: These research results indicate an evolutionary effect of HERVs and cell receptors and other proteins of the immune system and an allogeneic stimulation in pregnancy that is protective for life.

VII. About Testing, Responsibility and Ethics

What can be concluded from the findings mentioned above?

1. A HI-Virus has not been scientifically proven. It is gene expression in relation to the HLA-system for generating immune molecules in and between cells or a laboratory gene construct in experimental design.
2. People have differences in their MHC/HLA genes and their products; thus South Africans, naturally most often test HIV positive, which serves an evolutionary purpose.
3. To test HIV positive does not mean someone is infected with a virus.
4. A positive test result in mother and child is a normal biological process, as the mother fights the allograft (fetus) and the child receives the antibodies. Pregnancy is not a disease!
5. White blood cells have a genetic marker on the cell membrane that can stimulate allergic reactions through transfusion of blood products and by sexual contact with body fluids. Homosexuals and drug users have problems with allereactions (allergy) and stress but not with an exogenous virus. Other sexual transmitted diseases might be a problem, thus condoms are effective.
6. The HLA and HIV genes might be protective in other diseases (some cancers).
7. We have to discuss the terms “Virus” , “Exosomes” , “Endosomes” and “Genes” as evolution applies horizontal gene transfer and communication particles in normal biological processes like pregnancy and T1/T2-balance and in diseases [79] [80].
8. Stress might lead to an imbalance of the immune system and thus to disease.
9. Evolution is still going on. Thus we will find new “HIV-mutants” (transposition). The diversity of the MHC/HLA-system is granting co-evolution with microbes and is due to symbiogenesis.
10. We have to cease from testing because the tests have not been scientifically validated and provide false results. Testing for HIV is absolutely unethical!
11. Testing leads to fear and stress (**Nocebo Effect**) and might induce disease.
12. Treating with antiretrovirals might be harmful and even kill people. Drugs induce mutations and lead to disease. There is no specific disease that could be called AIDS. AIDS does not exist without testing. We have to be responsible in providing for the necessary living conditions of people worldwide, such as food, vitamins, selenium and minerals as well as pure drinking water, drug prevention and (oxidative) stress reduction.

- *We are responsible for what we do and what we neglect.*
- *Later generations will judge our behaviour in science and ethics.*
- *We must cease this misguided assessment immediately.*

References:

- [1] Novik, Nimrich et al: Epigenomics : Genome-Wide Study of Methylation Phenomena
Curr.Issues Mol. Biol (2002) b4 : 111-128
- [2] Jenuwein, Allis: Translating the Histone Code
Science 10 Aug. 2001: vol.293. no. 5532, pp. 1074-1080,
DOI : 10.1126/science.1063127
- [3] Van Lint, Emiliani, Verdin : Gene Expr. 1996 ; 5(4-5) : 245-53
- [4] Rakyan VK, Hildmann T, Novik KI, Lewin J, et al (2004)
DNA methylation profiling of the human major histocompatibility complex : A pilot
study for the human eiogenome project. PLOS Biol 2(12) : e405.
- [5] George Weinstock: Human Microbiome Project, Washington University in St. Louis,
School of Medicine
- [6] Samantha Martin, University of Liverpool: Scientists reveal driving force behind
Evolution, Nature, Samantha.martin@liv.ac.uk
- [7] Villareal, De Filippis: J.Virol. 74 7079 (2000), Lateral Gene Transfer or Viral
Colonisation?
- [8] Mooney, Cleland, Dep. Of Biol. Sciences, Stanford University. CA: The evolutionary
Impact of invasive species; www.pnas.org/cgi/doi/10.1073/pnas.091093398
- [9] Schuster, Miller, Ratan et.al; Nature **463**, 943-947 (18 Feb. 2010) doi: 10.1038/nat.08795
- [10] Bramanti, B.et al. Genetic discontinuity between local..., Science 326, 137-140
- [11] Malstrom, H. et al Ancient DNA reveals lack of continuity...
Curr.Biol.19, 1785-62 (2009)
- [12] Coop,G. et al. The role of geography in human adaption. PLoS Gen.5,e1000500 (2009)
- [13] Reich, D. et al. Reduced neutrophil count in people of African descent is due to ...
Gen.5,e1000360 (2009)
- [14] Hill, Allsopp et al. Extensive genetic diversity in the HLA class II region of Africans...
Institute of Molecular Medicine, Univ. of Oxford, John Radcliffe Hospital, Un. Kingd.
- [15] Kebede, Britton, Fehninger et al.Differences in Humoral Responses to the p 24 Antigen
Between Ethiopian and Swedish...; Clinical And Diagnostic Laboratory Immunology,
Sept. 1997,p.627-629 1071-412X/97\$04.00+0 Copyright 1997, Am. Soc. For Microb.
- [16] Mungall, Palmer et al.: The DNA sequence and analysis of human chromosome 6
The Wellcome Trust Sanger Institute, Genome Campus, Hinxton, Cambridge CB10
ISA, UK; Nature/Vol 425/ 23 OCTOBER 2003
- [17] Philip W. Hedrick: Evolutionary Genetics of the Major Histocompatibility Complex
The Americ. Naturalist, Vol. 143, No. 6 (Jun. 1994), pp. 945-964; The University of
Chicago Press; URL: <http://www.jstor.org/stable/2462890>
- [18] Gyllenstein, Sundvall et al.: Allelic diversity is generated by intraexon sequence ...
Proc. Of the Nat. Ac. Of Sci. US (1991), 88(9), 3686-90 ; PNASA6; ISSN: 0027-8424.
- [19] Doxiadis, de Groot et al. : Biomed. Prim. Res. Centre, Dep.of Comp. Gen. And
Refinement, The Netherl. ; Mol Immunol. 2008 May ; 45(10) :2743-51
- [20] C van Oosterhout : Population genomics and epidemiology
Transposons in the MHC...; Univ. of Hull, UK; Heredity(2009) 103, 190-191
- [21] John, Moore, James, Mallal, Centre for Clinical Immunology and Biomedical Statistics,
Royal Perth Hospital and Murdoch University, Western Australia, Program Abstr 8th
Conf Retrovir Oppor Infect Conf Retrovir Oppor Infect 8th 2001 Chic III. 2001 Feb 4-8:
93 (abstract no. 174).
- [21a] Trubey, Chertova et al.: J. of Virology, Dec. 2003, p. 12699-12709
- [21b] Louwagie, Janssens, Mascola et al. , J. of Virol., Jan 1995, p. 263-271 Vol 69, No. 1
Genetic diversity of the envelope gp from HIV Type 1 isolates of African Origin

- [22] Clerici, Shearer, *Immunology Today*, Vol. 15 No. 12 1994
- [23] Sriwanthana, Hodge, Mastro et al.: *AIDS Research and Human Retroviruses*. May 2001, 17(8): 719-734. doi: 10.1089/088922201750236997.
- [23a] Swann, Williams, Story et al., *Virology*, Volume 282, issue 2, 10 Apr. 2001, P. 267-277
- [24] Brass, Dyxhoorn et al.; *How. Hughes Med. Inst., Science* 2008 Feb 15; 319(5865):921-6
Identification of host proteins required for HIV infection through a functional genomic screen.
- [25] Brumme ZL, Brumme CJ, Heckermann D, et al. (2007) Evidence of diff. HLA class I mediated viral evolution... ; *PLoS Pathog* 3(7): e94. doi: 10.1371/journal.ppat.0030094
- [26] Shiina, Ota, Shimizu et al. *Genetics*, Vol. 173, 1555-1570, July 2006
doi : 10.1534/genetics.106.057034; Evolution of Major Histocompatibility Compl.
- [27] Avila-Rios, Ormsby et al., ; *Retrovirology*, 2009 Aug 10 ;6 :72. Unique features of HLA-mediated HIV evolution in a Mexican cohort : a comparative study.
- [28] Dalmaso, Carpentier et al.: (2008) The ANSR Genome Wide Association 01 Study. *PLoS ONE* 3(12): e3907. doi:10.1371/journal.pone.0003907
- [29] Dan Stowell “ HLA proteins are incorporated into budding HIV virus particles”
An article from “The Molecules of HIV”; www.mclld.co.uk/hiv
- [30] Shukla, Marques, Kimmel et al.: Human Chromosome 6-and 11-Encoded Factors Support HIV Type 1 Rev Function in A9 Cells ; *J. of Vir.*, Dec.1996, p. 9064-9068
- [31] Jeremias, Mockel, Witkin: Human semen induces interleukin 10 and 70 kDa heat shock protein gene transcription and ... *Molec. Hum. Reproduc.* vol 4 no11pp. 1084-88/1998
- [32] Slobedman, Barry, Spencer et al: Virus-Encoded Homologs of Cellular Interleukin-10 and Their Control of Host Immune Function; *J. of Virol.*, Oct. 2009, p.9618-29
doi: 10.1128/JVI.01098-09
- [33] Mathur, Goust, et al: Cross-reactivity of sperm and T lymphocyte antigens; *Am J Reprod Immunol.* 1981; 1(3) : 113-8
- [34] Pinto, Sharpe, Cohen, Shearer: Alloantigen-Stimulated Anti-HIV Activity; *Blood*, Vol.92 No. 9 (Nov.1), 1998: pp. 3346-3354
- [35] Padian, Shiboski, Glass, Vittinghoff: Heterosexual transmission of human immunodeficiency virus (HIV) in northern California: results from a ten-year study. *Am J Epidemiol.* 1997 Aug 15; 146(4):350-7.
- [36] Kingsley, PetersBabaahmady et al: Heterosexual and Homosexual Partners Practising Unprotected Sex May Develop Allogeneic Immunity and to a Lesser Extent Tolerance *PLoS ONE*, November 2009/ Volume 4/ issue 11/ e7938
- [37] Townsley-Fuchs, Kam, Fairhurst et al : Human immunodeficiency virus-1 (HIV-1) gp120 superantigen-binding serum antibodies. A host factor in homosexual HIV-1 transmission. *J Clin Invest.* 1996 Oct. 15; 98(8): 1794-1801. doi: 10.1172/ JCI1 18979.
- [38] Baccetti, Benedetto, Burrini et al: HIV-particles in spermatozoa of patients with AIDS and their transfer into the oocyte. *J Cell Biol.* 1994 Nov; 127(4): 903-14.
- [39] Calcaro PG, Whitmer KJ; International Conference on AIDS 1991 jun 16-21; 7: 62 (abstract no. TH.A.26). Does germ-line integration of HIV-1 result in expression during development?
- [40] Prescott, Taylor, Roper et al: Maternal reactivity to fetal alloantigens is related to Newborn immune responses and subsequent allergic disease. *Clinical and exp. Allergy* ISSN 0954-7894 Source 2005, vol. 35, n° 4, pp. 417-425 [9 page(s) (article)] (25ref.)
- [41] Arthos, Cicala, Martinelli, et al : HIV-1 envelope protein binds to and signals through Integrin $\alpha 4\beta 7$, the gut mucosal homing receptor for the peripheral T cells
Nature Immunology 9, 301-309 (2008) doi: 10. 1038/ni1566
- [42] Camus, Segura-Morales, Molle et al: The Clathrin Adaptor Complex AP-1 Binds HIV-1 and MLV Gag and Facilitates Their Budding.
Mol. Biol. Of the Cell Vol. 18, 3193-3203, August 2007

- [43] Teis, Saksena et al: ESCRT-II coordinates the assembly of ESCRT-III filaments for cargo sorting and multivesicular body vesicle formation. *The EMBO Journal* 29, 871-883 (03 March 2010) / doi: 10.1038/emboj.2009.408
- [44] Gumpert, Varco et al: Double-membrane gap junction internalization requires the clathrin-mediated endocytic machinery. *FEBS Lett.* 2008 Aug 20;582(19):2887-92.
- [45] Saurabh Mehandru, MD, Mount Sinai School of Medicine, and The Aaron Diamond AIDS Research Center, The Rockefeller University, New York: The Gastrointestinal Tract in HIV-1 Infection: Questions, Answers and More Questions! The PRN Notebook / Volume 12 / www.PRN.ORG December 2007 (online)
- [46] Stoorvogel, Kleijmeer et al: The Biogenesis and Function of Exosomes *Traffic.* 3(5): 321-330, May 2002
- [47] Gould, Booth, Hildreth: The Trojan exosome hypothesis 10592-10597 / *PNAS* / September 16, 2003 / vol. 100 / no. 19 www.pnas.org/cgi/doi/10.1073/pnas.1831413100
- [48] Frank P Ryan FRCP FLS: Human endogenous retroviruses in health and disease: a symbiotic perspective; *J of the Royal Society of Medicine* Vol. 97 Dec. 2004
- [49] Villareal LP.: Viruses and the Evolution of Life. Washington DC: Am Soc. For Microb.
- [50] Spruth, Stoiber et al: Neutralization of HIV Type 1 by Alloimmune Sera Derived from Polytransfused Patients. *AIDS Res. and Human Retrov.* April 1999, 15(6):533-543.
- [51] Goodglick, Zevit et al: Mapping the Ig superantigen-binding site of HIV-1 gp120 *The J. Immunol.* Vol.155, issue11 5151-5159, 1995
- [52] Woodland, Trudeau Institute: Immunity and retroviral superantigens in humans *Trends in Microbiology*, Vol. 23, Issue 2, 57-58, 1 February 2002
- [53] Hemalatha V, Skrikanth P, Mallika M. Superantigens – Concepts, clinical disease and therapy. *Indian J Med Microbiol* 2004;22:204-211
- [54] Omar, Hammond et al: HLA class I and II antigens in South African Blacks with Graves' disease (Univ. of Natal, Durban) doi:10.1016/0090-1229(90)90009-F (Elsevier)
- [55] Nishiyama, Karle, Planque et al: Antibodies to the superantigenic site of HIV-1 gp120: Hydrolytic and binding activities of the light chain subunit Online 12 January 2007
- [56] Talal, Garry, Schur et al: A conserved idiootype and antibodies to retroviral proteins in systemic lupus eryth. *J Clin. Invest.* 85(6): 1866-1871 (1990). **Doi:**10.1172/JCI114647.
- [57] E. Prem Kumar Reddy (Wistar Inst.in Philadelphia) in: The Associated Press- Jan. 1989: Similarity to AIDS multiple sclerosis tied to HTLV virus
- [58] Pavlova, Delev et al: Impact of polymorphisms of the major histocompatibility complex class II, interleukin-10, tumor necrosis factor-alpha and cytotoxic T-lymphocyte antigen-4 genes on inhibitor development in severe hemophilia A. *J Thromb Haemost.* 2009 Dec; 7(12):2006-2015.
- [59] Amman, Wara et al: Acquired Immunodeficiency in an Infant: Possible transmission by Means of Blood Products. *The Lancet*, Vol.321, Issue 8331, Ps. 956-958,30 April 1983 Doi: 10.1016/S0140-6736(83)92082-2
- [60] Baeten, Benki, Chohan et al: Hormonal contraceptive use, herpes simplex virus infection, and risk of HIV-1 acquisition among Kenyan women. *AIDS.* 2007 Aug 20;21(13):1771-7.
- [61] Alves, Meyer, Vieira, Toralles, Le Maire: Distribution and frequency of HLA alleles and Haplotypes in Brazilians with type 1 diabetes mellitus. *Arq Bras Endocrinol Metabol.* 2006 Jun; 50(3):436-44
- [62] Scanga, Bafica, Sher: Viral Gene Expression in HIV Transgenic Mice Is Activated by *Mycobacterium tuberculosis* and Suppressed after Antimycobacterial Chemotherapy *The J of Infec. Diseases* 2007; 195: 246-254 **DOI:** 10.1086/510244
- [63] Furth, Westphal, Henninghausen: Expression from the HIV-LTR is Stimulated by

- Glucocorticoids and Pregnancy. *AIDS Research and Human Retroviruses*. April 1990, 6(4): 553-560. doi: 10.1089/aid.1990.6.553.
- [64] Nonaka, Hashimoto et al: The human immunodeficiency virus type 1 Vpr protein and its carboxy-terminally truncated form induce apoptosis in tumor cells
Cancer Cell International 2009, 9:20doi:10.1186/1475-2867-9-20
- [65] Eleme, Taner et al: Cell Surface Organization of Stress-inducible Proteins ULBP and Mica That Stimulate Human NK Cells and T Cells via NKG2D
J. Exp. Med. The Rockefeller Univ. Press, Vol. 199, Number 7, Apr.5, 2004 1005-1010
- [66] Leserman, Petitto et al: Severe Stress, Depressive Symptoms, and Changes in Lymphocyte Subsets in Human Immunodeficiency Virus-Infected Men
Arch Gen Psychiatry. 1997; 54(3):279-285
- [67] Kovacsovics-Bankowski, Carini, Kashala, Essex ; *VIRAL IMMUNOLOGY*, 1992 Winter ;5(4) :243-8
Isotypic distribution of HIV-specific antibodies in individuals from Central Africa.
- [68] Hartigan, Simberkoff, Hamilton et al: Zidovudine resistance, syncytium-inducing phenotype, and HIV diseases progression in a case-control study
Journal of Acquired Immune Deficiency Syndromes ISSN: 0894-9255 / 1993 / Publisher: Lippincott Williams & Wilkins, WK Health
- [69] Hartley, Klasse, Sattentau, Moore: V3:HIV's switch-hitter
AIDS Res Hum Retroviruses. 2005 Feb;21(2):171-89.
- [70] Murray, De Koven, Spaner: Foreign Body Granuloma: A New Manifestation of Immune Restoration Syndrome / Online publication: 9 October 2002/ Presented at the 58th American Academy of Dermatology Annual Conference, San Francisco, California, March 2002 / *J Cutan Med Surg* 2003; 38-42 /DOI: 10.1007/s10227-002-2102-1
- [71] Reuse, Calao, Kabeya et al: Synergistic activation of HIV-1 expression by deacetylase inhibitors and prostratin: implications for treatment of latent infection.
PLoS One. 2009 Jun 30;4(6):e6093 / PMID: 2699633
- [72] Morcock, Thomas et al: HIV-1 inactivation by 4-vinylpyridine is enhanced by dissociating Zn²⁺ from nucleocapsid protein / Elsevier Inc. / online 4 March 2008 doi:10.1016/j.virol.2008.01.045
- [73] Dong Sung An, Yi-Ming Xie, Irvin S.Y.Chen: Envelope Gene of the Human Endogenous Retrovirus HERV-W Encodes a Functional Retrovirus Envelope
J Virol., Apr. 2001, p. 3488-3489. 2001 DOI: 10.1128/JVI.75.7.3488-3489.2001
- [74] Kumar, A. Kumar, S. Dinda, Luthra: Differential Expression of CXCR4 Receptor in Early and Term Human Placenta doi: 10.1016/j.placenta.2003.10.003 © Elsevier Ltd.
- [75] Gerlinskaya, Moshkin, Evsikov: Allogenic Stimulation in Early Pregnancy Improves Pre- and Postnatal Ontogenesis in BALB/cLac Mice
Journal of Reproduction and Development, Vol. 46, No. 6, 2000
- [76] Voiculescu, Avramescu et al / *Int Conf AIDS* 1993 Jun 6-11; 9:257 (abstr. No. PO-A-31-0737). HLA-DR/CD3+ cells in HIV-seropositive children at several stages of disease evolution.
- [77] Dilernia, Lourta, et al: Differential HLA-dependent HIV evolution among subtypes
Retrovir. 2006, 3(Suppl 1):P15doi:10.1186/1742-4690-3-S1-P15 Pub. 21. Dec.2006
- [78] Dan Stowell : HLA proteins are incorporated into budding HIV virus particles
An article from "The Molecules of HIV" / www.mclld.co.uk/hiv and Ott, D.E. (1997) Cellular Proteins in HIV Virions. Review in *Medical Virology* 7:167-180
- [79] AL Hughes: Origin and evolution of HLA class I pseudogenes / *Molecular Biology and Evolution*, Vol 12, 247-258, Copyright © 1995 by Soc. For Mol. Biol. And Evolution
- [80] Mattson, Taub: Ancient viral protein enrages astrocytes in multiple sclerosis
Nature Neuroscience 7, 1021-1023 (2004) doi :10.1038/nn1004-1021